



Oleaceae (tribus Oleeae)

Cartrema Raf.
Chionanthus L.
Forestiera Poir.
(*Fraxinus* L. → separate key)
Ligustrum L.
Nestegis Raf.
Notelaea Vent.
Olea L.
Osmanthus Lour.
Phillyrea L.
Picconia DC.
Syringa L.

VEGETATIVE KEY TO SPECIES CULTIVATED IN WESTERN EUROPE

Jan De Langhe

(8 March 2013 - 25 November 2020)

Vegetative identification key.

Introduction:

This key is based on vegetative characteristics, and therefore also of use when flowers and fruits are absent.

- Use a 10x hand lens to check pubescence, scales, teeth and venation pattern in general.
- Start counting veins at base of the lamina with first clearly ascending secondary vein, do not include intercalary veins, nor these ending in the apex.
- Look at the entire plant. Young specimens, shade, and strong shoots give an atypical view.
- Beware of hybridisation, especially with plants raised from seed other than wild origin.

Taxa treated in this key: → p10.

Taxa referred to synonymy in this key: → p10.

Remarks: → p10.

References:

- JDL herbarium and [illustrations](#)
- living specimens, in various arboreta, botanic gardens and collections
- literature:
 - Green, P.S. - (1997) - *Oleaceae* in The European Garden Flora VOL.5, p.574-592.
 - Grimshaw, J. & Bayton, R. - (2009) - *Oleaceae* in New Trees, 992p. AND [Trees and Shrubs Online](#)
 - Krüssmann, G. - (1976) - *Chionanthus* in Handbuch der Laubgehölze, VOL 1 p.349-350.
 - Krüssmann, G. - (1977) - *Forestiera, Ligustrum, Notelaea, Olea, Osmanthus* and *Phillyrea* in Handbuch der Laubgehölze, VOL 2 466p.
 - Krüssmann, G. - (1978) - *Syringa* in Handbuch der Laubgehölze, VOL 3 p.395-408.
 - Rehder, A. - (1940) - *Oleaceae* in Manual of cultivated trees and shrubs hardy in North America, p.765-793.
 - Lancaster, R. - (2019) - *Oleaceae* in The Hillier Manual of Trees & Shrubs 9yh Edition, 637p.
 - Royal Botanic Gardens Kew. The genus *Ligustrum* in World checklist of Selected Plantfamilies [online](#)
 - Royal Botanic Gardens Kew. The genus *Syringa* in World checklist of Selected Plantfamilies [online](#)
 - Chang, M.-C., Qiu, L.-O., Green, P.S. - (1996) - *Oleaceae* in Flora of China VOL.15, p.272-319. [online edition](#)

Acknowledgements:

I am particularly grateful to Wolfgang Bopp, Peter Brownless, Barry Clarke, Koen Camelbeke, Gert Fortgens, Francisco Garin, Paul Goetghebeur, Tom Hudson, Jean Merret, Jef Van Meulder, Joke Osaer and Rolf Zumbrunn for extra help with constructive comments and specimens.

My explicit thanks for support goes to Arboretum Bokrijk, Arboretum Keracoual, Arboretum Trompenburg Rotterdam, Arboretum Wespelaar, Ghent University Botanical Garden, Hillier Gardens and Arboretum, Iturraran Botanical Garden, Jardin Botanique de l'Université de Strasbourg, Royal Botanic Gardens Edinburgh, Tregrehan Garden and Trompenburg Tuinen en Arboretum.

Also thanks to the responsible persons of various arboreta, botanic gardens and collections where I could collect herbarium specimens since 1982.

Copyright © 2013-2020 **Jan De Langhe**

Non-commercial use authorized with mentioning "Jan De Langhe - Ghent University Botanical Garden".

[Plantentuin Universiteit Gent](#)

[Arboretum Wespelaar](#)

KEY TO GROUPS

- 01 a Leaf at least in part compound (with 3-more leaflets), OR/IF simple then with lamina pinnatisect to deeply lobed (lobe length >1/3 lamina width). **GROUP A → p2**
- b Leaf simple. Lamina margin entire, dentate or (sub)lobed (lobe length <1/3 lamina width). **02**
- 02 a Lamina margin often dentate: varying from minutely crenulate to serrulate or distantly dentate to (sub)lobed (teeth and lobe height/length <1/3 lamina width). **03**
- b Lamina margin predominantly entire (rarely few shoots with minutely dentate leaves). . **04**
- 03 a Dentate/lobed lamina margin with teeth spinulose. **GROUP B1 → p3**
- b Dentate/lobed lamina margin with teeth acute to obtuse. **GROUP B2 → p3**
- 04 a Lamina LS finely reticulate: tertiary vein areoles and sometimes up to quaternary vein areoles quite visible (10x LENS). **05**
- b Lamina LS poorly reticulate: tertiary vein areoles not or partly visible. **07**
- 05 a Terminal bud abortive, reduced, or replaced by 2 lateral buds. **GROUP C1 → p5**
- b Terminal bud predominantly present. **06**
- 06 a Terminal bud with 2 (sub)valvate scales visible (10x LENS). **GROUP C2 → p5**
- b Terminal bud with several imbricate scales visible. **GROUP C3 → p6**
- 07 a Petiole predominantly 6-10(-20) mm. **GROUP D1 → p7**
- b Petiole predominantly 2-6 mm. **GROUP D2 → p8**

GROUP A

LEAF at least in part COMPOUND (with 3-more leaflets),
OR/IF SIMPLE then lamina pinnatisect to deeply lobed (lobe length >1/3 lamina width).

- 01 a Leaf pinnately compound. **02**
- b Leaf simple, with lamina variable: both entire AND variable deeply lobed to pinnatisect. **03**
- 02 a Lamina with 7-11 leaflets AND leaflet sessile with margin entire. ***Syringa pinnatifolia***
Lamina not simultaneously with the characters mentioned above. → genus *Fraxinus*
- 03 a Pinnatisect leaves with 2-4 narrow lobes/side. ***Syringa protolaciniata***
- b Pinnatisect leaves with ≤3 narrow lobes/side. **04**
- 04 a Pinnatisect leaves with 1-3 narrow lobes/side. ***Syringa ×laciñata***
- b Pinnatisect leaves with 1-2 narrow lobes/side. ***Syringa ×diversifolia***

GROUP B1
LAMINA DENTATE/LOBED - margin with teeth spinulose -

- 01 a Lamina LS reticulation rather strongly developed: quaternary vein areoles between two secondary veins numerous and obvious (10x LENS). 02
 - b Lamina LS reticulation rather weakly developed: tertiary vein areoles between two secondary veins few or partly obscured. 03
- 02 a Dentate lamina with forward directed small teeth, ≤ 2 mm. *Osmanthus fragrans*
 - b Dentate lamina with sideward directed larger teeth, 2-4 mm. *Osmanthus yunnanensis*
- 03 a Petiole predominantly ≤ 6 mm. 04
 - b Petiole predominantly > 6 mm. 06
- 04 a Dentate lamina margin with > 10 teeth/side. *Phillyrea latifolia*
 - b Dentate lamina margin predominantly with ≤ 10 teeth/side. 05
- 05 a Lamina rigid and texture very thick, US and LS secondary veins obscure and only visible in part with 10x LENS. *Osmanthus delavayi*
 - b Lamina +/- supple and texture leathery, US and LS secondary veins +/- visible and quite so with 10x LENS. *Osmanthus delavayi* 'Latifolius'
- 06 a Dentate lamina margin minutely spinulose dentate, teeth ≤ 2 mm. .. *Osmanthus serrulatus*
 - b Dentate lamina margin coarsely spinulose dentate to lobed, teeth (lobes) 3-10 mm. 07
- 07 a Dentate lamina margin with 1-3 large +/- triangular teeth-lobes/side.
 - *Osmanthus heterophyllus*
 - b Dentate lamina margin with (2-)4-10(-12) coarse teeth-lobes/side. 08
- 08 a Lamina length/width predominantly $\geq 2/1$, largest teeth > 5 mm. *Osmanthus armatus*
 - b Lamina length/width predominantly $\leq 2/1$, largest teeth < 5 mm. *Osmanthus ×fortunei*

GROUP B2
LAMINA DENTATE/LOBED - margin with teeth acute to obtuse -

- 01 a Shoot at least in part with lateral buds in whorls of three.
 - *Fraxinus angustifolia* 'Monophylla'
 - b Shoot with lateral buds (sub)opposite. 02
- 02 a Current year shoot 4-ribbed. Secondary year shoot often with internodes 4-ribbed.
 - *Fraxinus anomala*
 - b Current year shoot flattened and/or widened at nodes. 03
- 03 a Terminal bud 5-10 mm AND black. *Fraxinus excelsior* 'Diversifolia'
 - b Terminal bud < 5 mm AND/OR brown to greenish. 04

04 a Lamina L/W 6/1-4/1.	<i>Phillyrea angustifolia</i>
b Lamina L/W ≤4/1.	05
05 a Lamina midvein length predominantly 1-4(-5) cm <u>AND</u> margin with 6-10(-12) teeth/side.	06
b Lamina midvein length predominantly >4 cm, <u>OR</u> margin predominantly with 10-20 teeth/side.	08
06 a Lamina narrowly elliptic to oblanceolate, base attenuate. Margin entire to minutely dentate with inward directed teeth.	<i>Forestiera neomexicana</i>
b Lamina broadly elliptic to (sub)orbicular, base broadly cuneate to rounded. Margin dentate with outward directed teeth.	07
07 a Lamina rigid and texture very thick, US and LS secondary veins obscure and only visible in part with 10x LENS.	<i>Osmanthus delavayi</i>
b Lamina +/- supple and texture leathery, US and LS secondary veins +/- visible and quite so with 10x LENS.	<i>Osmanthus delavayi</i> 'Latifolius'
08 a Lamina apex predominantly obtuse to rounded.	09
b Lamina apex acute to (short) acuminate.	11
09 a Plant deciduous. Lamina LS venation pubescent, with up to quaternary vein areoles clearly visible (10x LENS).	<i>Chionanthus retusus</i>
b Plant evergreen. Lamina LS venation glabrous, with tertiary vein areoles not or slightly visible in part.	10
10 a Dentate lamina margin with teeth restricted to apical half. Petiole predominantly 6-10 mm.	<i>Picconia azorica</i>
b Dentate lamina margin with teeth often from base to apex. Petiole predominantly 2-6 mm.	<i>Phillyrea latifolia</i>
11 a Lamina midvein length 7-15 cm.	<i>Picconia excelsa</i>
b Lamina midvein length <8 cm.	12
12 a Lamina margin with 20-50 teeth/side.	13
b Lamina margin with (10-)15-20 teeth/side.	14
13 a Lamina LS glabrous.	<i>Osmanthus suavis</i>
b Lamina LS midvein pubescent and surface scattered pubescent.	<i>Forestiera ligustrina</i>
14 a Lamina apex obtuse to acute.	<i>Phillyrea latifolia</i>
b Lamina apex acute to (sub)acuminate.	15
15 a Lamina LS irregularly mottled and dotted (10x LENS).	<i>Osmanthus × burkwoodii</i>
b Lamina LS regularly and very finely dotted .	<i>Osmanthus delavayi</i> × <i>suavis</i>

GROUP C1
LAMINA ENTIRE WITH LS FINELY RETICULATE (10 \times LENS).
Terminal bud abortive, reduced, or replaced by 2 buds.

- 01 a Lamina base predominantly (sub)cordate to (sub)truncate 02
 - b Lamina base predominantly rounded to cuneate 03
- 02 a Lamina ovate, apex rather gradually long acuminate [flowers mid-May]. .. ***Syringa vulgaris***
 - b Lamina broadly ovate to orbicular-ovate, apex rather acute to abruptly acuminate [flowers early May]. ***Syringa oblata***
 - Hybrids between previous taxa flowering from mid-April. ***Syringa ×hyacinthiflora***
- 03 a Lamina width often >6 cm. ***Syringa reticulata***
 - Lamina LS pubescent at least on midvein. ***Syringa reticulata* subsp. *reticulata***
 - Lamina LS glabrous. ***Syringa reticulata* subsp. *amurensis***
- b Lamina width rarely >6 cm. 04
- 04 a Lamina broadly ovate to (sub)orbicular. ***Syringa meyeri***
 - b Lamina ovate. 05
- 05 a Lamina LS clearly pubescent AND secondary veins clearly elevated (10 \times LENS).
 - ***Syringa pubescens***
 - Shoot pubescent, lamina LS venation pubescent. ***Syringa pubescens* subsp. *microphylla***
 - Shoot glabrous to glabrescent, lamina LS secondary veins +/- pubescent. ***Syringa pubescens* subsp. *patula***
- b Lamina LS glabrous to glabrescent AND secondary veins not or (slightly) elevated in part. 06
- 06 a Lamina LS glabrous to glabrescent AND secondary veins not to slightly elevated (10 \times LENS). ***Syringa reticulata* subsp. *pekinensis***
 - b Lamina LS glabrous AND larger secondary veins clearly elevated. ***Syringa ×chinensis***

GROUP C2
LAMINA ENTIRE WITH LS FINELY RETICULATE (10 \times LENS).
Terminal bud scales valvate.

- 02 a Lamina lanceolate: L/W \geq 6/1, texture thick leathery. ***Nestegis cunninghamii***
 - b Lamina elliptic, ovate or obovate: L/W \leq 4/1, texture +/- papery and thin. 03
- 03 a Petiole \leq 10 mm AND lamina predominantly 3-10 cm. ***Chionanthus pygmaeus***
 - b Petiole 10-30 mm, OR lamina often 10-20 cm. 04

- 04 a Lamina apex obtuse to rounded, rarely acute AND largest leaves with secondary veins ≤ 7 /side (intercalary veins NOT included). *Chionanthus retusus*
- b Lamina apex acute to acuminate AND largest leaves with secondary veins 7-10-(more)/side (intercalary veins NOT included). *Chionanthus virginicus*

GROUP C3
LAMINA ENTIRE WITH LS FINELY RETICULATE (10 \times LENS).
 Current year shoot terminal bud scales imbricate.

- 01 a Lamina L/W $\geq 3/1$ AND base attenuate. *Forestiera neomexicana*
- b Lamina L/W $\leq 3/1$, OR base cuneate, or rounded to cordate. 02
- 02 a Plant evergreen. Lamina +/- thick leathery. 03
- b Plant deciduous. Lamina thin, papery. 04
- 03 a Lamina margin undulate in most leaves. *Osmanthus yunnanensis*
- b Lamina margin straight. *Osmanthus decorus*
- 04 a Shoot slender: $\emptyset \leq 2$ mm AND lamina margin fimbriate (10 \times LENS). *Syringa tomentella*
- Lamina LS (almost) glabrous (10 \times LENS). *Syringa tomentella* subsp. *yunnanensis*
 - Lamina LS clearly pubescent.
 - Lamina US pubescent, LS pubescent. *Syringa tomentella* subsp. *tomentella*
 - Lamina US glabrous, LS midvein and secondary veins pubescent. *Syringa tomentella* subsp. *sweginzowii*
- b Shoot +/- firm: $\emptyset > 2$ mm, AND/OR lamina margin +/- ciliate. 05
- 05 a Lamina LS papillose AND glabrous (10 \times LENS). *Syringa emodi*
- b Lamina LS not papillose, OR pubescent (10 \times LENS). 06
- 06 a Lamina midvein length 8-18 cm AND apex acute to acuminate or acuminate to caudate in part of the leaves [panicle nodding or pendulous]. *Syringa komarovii*
- Panicle rather cylindrical, compact, $\emptyset \leq 5$ cm. *Syringa komarovii* subsp. *komarovii*
 - Panicle rather cylindrical-pyramidal, often interrupted, $\emptyset 3-10$ cm. *Syringa komarovii* subsp. *reflexa*
- b Lamina apex acute to acuminate (never caudate), OR midvein length predominantly 5-12 cm [panicle +/- erect or spreading]. 07
- 08 a Lamina predominantly: broadly elliptic, LS glaucous/whitish green AND midvein length 5-12(-14) cm [panicle ≤ 20 cm; corolla tube funnel-shaped, lobes erect]. *Syringa josikaea*
- b Lamina predominantly broadly elliptic to oblong, OR LS pale green OR midvein length 8-18 cm [panicle > 20 cm, OR corolla cylindrical, OR lobes spreading]. *Syringa villosa*
- Shoot soon pale grey. Lamina LS rather pale green [panicle 20-30 cm; corolla tube funnel-shaped, lobes erect, tip incurved].... *Syringa villosa* subsp. *wolfii*
 - Shoot soon brownish. Lamina LS rather glaucous green [panicle < 20 cm; corolla tube cylindrical, lobes spreading]. *Syringa villosa* subsp. *villosa*

GROUP D1
LAMINA ENTIRE WITH LS POORLY RETICULATE (10x LENS).
Petiole predominantly 6-10(-20) mm.

- 01 a Plant pubescent on lamina LS or petiole or shoot. 02
 - b Plant glabrous. 03
- 02 a Lamina LS secondary veins often 8-10/side. *Ligustrum sinense*
 - b Lamina LS secondary veins predominantly 6-8/side. *Ligustrum foliosum*
- 03 a Lamina apex predominantly obtuse to rounded. *Picconia azorica*
 - b Lamina apex acute to (short) acuminate pr +/- caudate. 04
- 04 a Lamina thin papery. 05
 - b Lamina +/- thick leathery. 06
- 05 a Secondary veins (6-)10-16/side. Apex acuminate to +/- caudate. *Ligustrum compactum*
 - b Secondary veins 6-10/side. Apex acute to acuminate. *Ligustrum sinense*
- 06 a Lamina lanceolate to narrowly elliptic L/W predominantly >4/1. *Nestegis lanceolata*
 - b Lamina ovate to elliptic L/W predominantly <4/1. 07
- 07 a Lamina often wavy AND/OR margin often undulate. 08
 - b Lamina margin rather straight or almost so. 09
- 08 a Lamina LS secondary veins anastomosing near margin (10x LENS). *Picconia excelsa*
 - b Lamina LS secondary veins forming areoles near margin. *Nestegis apetala*
- 09 a Lamina <4/1 AND obovate to oblanceolate, or (narrowly) elliptic to oblanceolate AND base attenuate. *Cartrema americana*
 - b Lamina ≤3/1, OR ovate to elliptic or elliptic to oblong, or oblong to linear, OR base rounded to (broadly) cuneate. 10
- 10 a Lamina US venation obscure to the naked eye. 11
 - b Lamina US venation clearly visible. 12
- 11 a Lamina apex obtuse to acute, secondary veins ≤6/side. *Ligustrum japonicum*
 - b Lamina apex acute to acuminate, secondary veins 6-8/side. *Ligustrum lucidum*
- 12 a Lamina apex acute to acuminate. *Osmanthus serrulatus*
 - b Lamina apex acute. 13
- 13 a Lamina LS several tertiary veins clearly visible between each secondary vein (10x LENS). *Osmanthus decorus*
 - b Lamina LS tertiary veins obscured to only slightly visible between some secondary veins usually near margin. 13

- 13 a Spring flowering evergreen shrub. *Osmanthus marginatus*
 b Autumn flowering evergreen shrub. *Osmanthus rigidus*

GROUP D2
 LAMINA ENTIRE WITH LS POORLY RETICULATE (10x LENS).
 Petiole predominantly 2-6 mm.

- 01 a Lamina LS, at least in part, scattered to very densely grey-silvery scaly (10x LENS!!! sometimes it is necessary to check under different angels). *Olea europaea*
 b Lamina LS not scaly, often minutely (usually dark) dotted. 02
- 02 a Secondary vein axils often with domatia (10x LENS). *Chionanthus foveolatus*
 b Secondary vein axils without domatia. 03
- 03 a Shoot with two or more serial buds often present. 04
 b Shoot with serial buds absent. 06
- 04 a Lamina LS midvein depressed. *Notelaea ligustrina*
 b Lamina LS midvein elevated. 05
- 05 a Lamina length/width 6/1-4/1, base cuneate. *Phillyrea angustifolia*
 b Lamina length/width 3/1-2/1, base broadly cuneate to rounded or (sub)cordate. *Phillyrea latifolia*
- 06 a Lamina texture thick and rigid, 15-50 × 12-30 mm (larger on vigorous plants/shoots). *Ligustrum sempervirens*
 b Lamina texture rather thin and supple. 07
- 07 a Lamina size predominantly 10-30(-40) × 10-15(-20) mm. 08
 b Lamina size predominantly ≥40 × 20 mm. 11
- 08 a Lamina broadly ovate with largest width in or below the middle. *Ligustrum henryi*
 b Lamina broadly elliptic to (sub)orbicular with largest width in or sometimes above the middle. 09
- 09 a Lamina apex predominantly emarginate. *Ligustrum sinense*
 b Lamina apex acute to obtuse. 10
- 10 a Lamina LS with ≥3 rather obvious secondary veins/side. *Ligustrum delavayanum*
 b Lamina with 2(-3) rather obscure secondary veins/side. *Ligustrum morrisonense*
- 11 a Lamina apex predominantly obtuse to rounded, sometimes +/- retuse. 12
 b Lamina apex predominantly acute to (short) acuminate, OR very variable: shoots with lamina apex acute and shoots with lamina apex obtuse or shoots with both kinds. 14

- 12 a Lamina base predominantly attenuate. *Ligustrum quihoui*
- b Lamina base predominantly rounded to cuneate. 13
- 13 a Petiole predominantly 1-3 mm. *Ligustrum obtusifolium*
- b Petiole 2-6(-more) mm. *Ligustrum sinensis*
- 14 a Lamina length/width 5/1-10/1. *Nestegis montana*
- a Lamina L/W 4/1. 15
- 15 a Plant glabrescent to glabrous (10x LENS). 16
- b Plant clearly pubescent on shoot or lamina LS midvein and margin. 17
- 16 a Lamina (narrowly) ovate, with L/W 2/1-3/1. *Ligustrum ovalifolium*
- b Lamina narrowly ovate to lanceolate, with L/W 2/1-4/1. *Ligustrum vulgare*
- 17 a Lamina apex acute to (short) acuminate (petiole 3-5 mm). *Ligustrum tschonoskii*
- b Lamina apex acute to obtuse. 18
- 18 a Petiole 3-6 mm. *Ligustrum sinense*
- b Petiole predominantly <3 mm. *Ligustrum obtusifolium*
- Hybrid with *L. ovalifolium*: Lamina apex very variable: shoots with lamina apex acute, others with apex obtuse and shoots with both kinds of lamina present. *Ligustrum ×ibolium*

Taxa treated in this key.

<i>Cartrema americana</i>	<i>Osmanthus ×fortunei</i> (<i>O. fragrans</i> × <i>O. heterophyllus</i>)
<i>Chionanthus foveolatus</i>	<i>Osmanthus fragrans</i>
<i>Chionanthus pygmaeus</i>	<i>Osmanthus heterophyllus</i>
<i>Chionanthus retusus</i>	<i>Osmanthus marginatus</i>
<i>Chionanthus virginicus</i>	<i>Osmanthus rigidus</i>
<i>Forestiera ligustrina</i>	<i>Osmanthus serrulatus</i>
<i>Forestiera neomexicana</i>	<i>Osmanthus suavis</i>
<i>Ligustrum compactum</i>	<i>Osmanthus yunnanensis</i>
<i>Ligustrum delavayianum</i>	
<i>Ligustrum foliosum</i>	
<i>Ligustrum henryi</i>	
<i>Ligustrum ×ibolium</i> (<i>L. obtusifolium</i> × <i>L. ovalifolium</i>)	
<i>Ligustrum japonicum</i>	
<i>Ligustrum lucidum</i>	
<i>Ligustrum morissonense</i>	
<i>Ligustrum obtusifolium</i>	
<i>Ligustrum ovalifolium</i>	
<i>Ligustrum quihoui</i>	
<i>Ligustrum sempervirens</i>	
<i>Ligustrum sinense</i>	
<i>Ligustrum tschonoskii</i>	
<i>Ligustrum vulgare</i>	
<i>Nestegis apetala</i>	
<i>Nestegis cunninghamii</i>	
<i>Nestegis montana</i>	
<i>Nestegis lanceolata</i>	
<i>Notelaea ligustrina</i>	
<i>Olea europaea</i>	
<i>Osmanthus americanus</i>	
<i>Osmanthus armatus</i>	
<i>Osmanthus ×burkwoodii</i> (<i>O. delavayi</i> × <i>O. decorus</i>)	
<i>Osmanthus decorus</i>	
<i>Osmanthus delavayi</i>	
<i>Osmanthus delavayi</i> 'Latifolius'	
<i>Osmanthus delavayi</i> × <i>suavis</i>	
	<i>Phillyrea angustifolia</i>
	<i>Phillyrea latifolia</i>
	<i>Picconia azorica</i>
	<i>Picconia excelsa</i>
	<i>Syringa emodi</i>
	<i>Syringa josikaea</i>
	<i>Syringa komarovii</i>
	- subsp. <i>komarovii</i>
	- subsp. <i>reflexa</i>
	<i>Syringa meyeri</i>
	<i>Syringa oblata</i>
	<i>Syringa pinnatifolia</i>
	<i>Syringa protolaciniata</i>
	<i>Syringa pubescens</i>
	- subsp. <i>microphylla</i>
	- subsp. <i>patula</i>
	- subsp. <i>pubescens</i>
	<i>Syringa reticulata</i>
	- subsp. <i>amurensis</i>
	- subsp. <i>pekinensis</i>
	- subsp. <i>reticulata</i>
	<i>Syringa tomentella</i>
	- subsp. <i>sweginzowii</i>
	- subsp. <i>tomentella</i>
	- subsp. <i>yunnanensis</i>
	<i>Syringa villosa</i>
	- subsp. <i>villosa</i>
	- subsp. <i>wolfii</i>
	<i>Syringa vulgaris</i>

Taxa referred to synonymy in this key.

<i>Ligustrum amurense</i> → <i>L. obtusifolium</i> var. <i>suave</i>	<i>Syringa chinensis</i> → <i>S. oblata</i>
<i>Ligustrum chenaultii</i> → <i>L. compactum</i>	<i>Syringa fauriei</i> → <i>S. reticulata</i> subsp. <i>amurensis</i>
<i>Ligustrum confusum</i> Hort. → <i>L. compactum</i>	<i>Syringa giraldiana</i> → <i>S. pubescens</i> subsp. <i>microphylla</i>
<i>Ligustrum indicum</i> → <i>L. sinense</i>	<i>Syringa japonica</i> → <i>S. reticulata</i> subsp. <i>reticulata</i>
<i>Ligustrum insulare</i> → <i>L. vulgare</i>	<i>Syringa julianae</i> → <i>S. pubescens</i> subsp. <i>microphylla</i>
<i>Ligustrum microcarpum</i> → <i>L. sinense</i>	<i>Syringa microphylla</i> → <i>S. pubescens</i> subsp. <i>microphylla</i>
<i>Ligustrum stauntonii</i> → <i>L. sinense</i>	<i>Syringa reflexa</i> → <i>S. komarovii</i> subsp. <i>reflexa</i>
<i>Osmanthus americanus</i> → <i>Cartrema americana</i>	<i>Syringa pekinensis</i> → <i>S. reticulata</i> subsp. <i>pekinensis</i>
<i>Osmanthus forrestii</i> → <i>O. yunnanensis</i>	<i>Syringa patula</i> → <i>S. pubescens</i> subsp. <i>patula</i>
<i>Osmanthus omeiensis</i> → <i>O. marginatus</i>	<i>Syringa potaninii</i> → <i>S. pubescens</i> subsp. <i>microphylla</i>
<i>Phillyrea media</i> → <i>P. latifolia</i>	<i>Syringa sweginzowii</i> → <i>S. tomentella</i> subsp. <i>sweginzowii</i>
<i>Syringa afganica</i> → <i>S. protolaciniata</i>	<i>Syringa tigerstedtii</i> → <i>S. tomentella</i> subsp. <i>sweginzowii</i>
<i>Syringa amurensis</i> → <i>S. reticulata</i> subsp. <i>amurensis</i>	<i>Syringa velutina</i> → <i>S. pubescens</i> subsp. <i>patula</i>
	<i>Syringa yunnanensis</i> → <i>S. tomentella</i> subsp. <i>yunnanensis</i>
	<i>Syringa wolfii</i> → <i>S. villosa</i> subsp. <i>wolfii</i>

Remarks.

Plants in collections labelled *L. confusum*, *L. ibota*, *L. leucanthum*, *L. lindleyi*, *L. salicinum* are rather sporadically encountered. These are excluded in this key for the time being.

Ligustrum confusum with large leaves (lamina often >10 cm long) is *L. compactum*.

L. strongylophyllum with large leaves (lamina 5-8 × 2-3 cm with an obtuse to rounded apex) is usually *L. obtusifolium*.

