



Fabaceae (pro parte)

Acacia Mill.

Albizia Durazz.

Biancaea Tod.

Ceratonia L.

Cercis L.

Cladrastis Raf.

Erythrostemon Klotzsch

Gleditsia L.

Gymnocladus Lam.

+*Laburnocytisus* C.K.Schneid.

Laburnum Fabr.

Maackia Rupr.

Paraserianthes I.C.Nielsen

Platyosprion Maxim.

Robinia L.

Sophora L.

Styphnolobium Schott

Tipuana Benth.

Vachellia Wight & Arn.

VEGETATIVE KEY TO ARBORESCENT SPECIES CULTIVATED IN WESTERN EUROPE

Jan De Langhe

(31 August 2011 - 12 September 2023)

Vegetative identification key.

Introduction:

The taxa discussed here are arborescent (in a few cases exceptions are made to complement an otherwise arborescent genus). The *Albizia*, *Biancaea*, *Erythrostemon*, *Tipuana* and selected *Acacia* taxa proved to be hardy since several decades in the Mediterranean region and more and more also beyond. It are popular container plants, often cultivated under glass.

This key is based on vegetative characteristics, and therefore also of use when flowers and fruits are absent.

- Use a 10x hand lens to check pubescence and venation pattern in general.
- Look at the entire plant. Young specimens, shade, and strong shoots give an atypical view, so avoid them!
- Beware of hybridisation, especially with plants raised from seed other than wild origin.

Abbreviations and glossary for this key:

LS → lower surface

L/W ratio → length/width ratio

US → upper surface

jugary gland → gland on the rachis of a pinnate leaf near the attachment of pinnae (jugary extra-floral nectary)

phyllode → modified petiole, with the appearance and function of a leaf

pinna (pinnae) → primary segment(s) of a bipinnate leaf

pulvinus → enlarged/thickened basal part of a petiole

stipel (stipels) → stipule(s) of a leaflet

stipule (stipules) → leaflike structure(s) at the base of a leaf or its petiole

stipular spine → sharp-pointed stipule

Taxa treated in this key: → **p9**.

Taxa referred to synonymy in this key: → **p9**.

References:

- JDL herbarium and [illustrations](#)
- living specimens, in various arboreta, botanic gardens and collections
- literature:

Flora of China - (2010) - **Fabaceae** in Flora of China VOL.10, p.1-577. [online edition](#)

Grimshaw, J. & Christian, T. & all - **Fabaceae** in [Trees and Shrubs Online](#)

Jacquemin, D. - (1997) - Mimosas pour le Climat Méditerranéen – Une étude du genre *Acacia*, 128p.

Krüssmann, G. - (1976-1978) - **Fabaceae** in Handbuch der Laubgehölze, 3VOL, 1585p.

Lancaster, R. - (2019) - **Fabaceae** in The Hillier Manual of Trees & Shrubs 9th Edition, 637p.

Maslin, B.R. (coordinator) - (2018) - **WATTLE**, Interactive Identification of Australian *Acacia*. Version 3. [WATTLE](#)

Acknowledgements:

I am particularly grateful to Benn Baudts, Kenneth Bauters, Wolfgang Bopp, Koen Camelbeke, Tom Christian, Allen Coombes, Dan Crowley, Dirk De Meyere, Siegfried Gand, Paul Goetghebeur, John Grimshaw, Tom Hart Dyke, Tom Hudson, David Jewell, Owen Johnson, Raf Lenaerts, Conny Löhne, Bruce Maslin, Daniel Murphy, Ralf Omlor, Joke Ossaer, Frédéric Tournay, Liesbeth Uijtewaal, Filip Van den Bossche, Marian Vande Wiele, Jef Van Meulder, Kurt Van Nieuwenhuyse, Jan Willem Vos, Neville Walsh, Maximilian Weigend, Laura Wester, Johan Willm, Erik Windey, and Rolf Zumbunn for extra help with constructive comments and specimens.

My explicit thanks for support goes to Arboretum Bokrijk, Arboretum Het Leen, Arboretum Wespelaar, Botanischer Gärten der Universität Bern, Botanischer Gärten der Universität Bonn, Botanischer Garten der Universität Zürich, Botanischer Garten Johannes Gutenberg-Universität Mainz, Botanischer Garten Köln, Hortus Botanicus Amsterdam Buitenveldert, Jardin Botanique de la Villa Thuret, Jardin Botanique Université de Fribourg, Lullingstone Castle & The World Garden, Merrigum, Parc Botanique Château Pérouse, Plantentuin Meise, Plantentuin Universiteit Gent, Royal Botanic Gardens Kew, Sir Harold Hillier Gardens, Tregrehan Garden, Tresco Abbey Gardens, and Westonbirt Arboretum.

An extra word of thanks goes to Bruce Maslin, for his professionalism and his patience in answering all my questions.

Thanks also to the numerous managers of arboreta, botanical gardens and collections where I have been able to take herbarium specimens since 1982.

Copyright © 2011-2023 **Jan De Langhe**

Non-commercial use authorized with mentioning "Jan De Langhe - Ghent University Botanical Garden".

[Plantentuin Universiteit Gent](#)

[Arboretum Wespelaar](#)

KEY TO GROUPS

- 01 a **Foliage with simple leaves, OR with phyllodes (in the case of the latter sometimes juvenile bipinnate leaves may occur in part of the foliage).** 02
- b **Foliage with all leaves compound (trifoliolate, pinnate or bipinnate).** 03
- 02 a **Foliage with simple leaves.**
Group A1: *Cercis*. →page 3
- b **Foliage with phyllodes.**
Group A2: *Acacia* (pro parte). →page 4
- 03 a **Leaves trifoliolate.**
Group B1: +*Laburnocytisus*, *Laburnum*. →page 5
- b **Leaves pinnate AND/OR bipinnate.** 04
- 04 a **Leaflet margin (sometimes minutely!) crenulate (10x LENS, examine several leaves).**
Group B2: *Gleditsia*. →page 5
- b **Leaflet margin entire.** 05
- 05 a **Leaves bipinnate.**
Group B3: *Acacia* (pro parte), *Albizia*, *Biancaea*, *Erythrostemon*, *Gymnocladus*, *Paraserianthes*, *Vachellia*. →page 6
- b **Leaves pinnate.** 06
- 06 a **Lateral buds concealed by leaf pulvini.**
Group B4: *Cladrastis*, *Platyosprion*, *Robinia*, *Styphnolobium*. →page 7
- b **Lateral buds visible.**
Group B5: *Ceratonia*, *Maackia*, *Sophora*, *Tipuana*. →page 8

**GROUP A1:
Foliage with simple leaves.**

- 01 a Lamina apex predominantly obtuse AND/OR emarginate. 02
 b Lamina apex predominantly acute to acuminate. 05
- 02 a Juvenile branchlet, petiole and lamina lower surface venation pubescent (10x LENS).
 ***Cercis canadensis* var. *mexicana***
- b Juvenile branchlet, petiole and lamina lower surface glabrous (except vein axils). 03
- 03 a Lamina predominantly reniform. ***Cercis griffithii***
 b Lamina orbicular, OR orbicular to reniform. 04
- 04 a Tree. Lamina texture thin/papery. ***Cercis siliquastrum***
 b Spreading multi-stemmed shrub, rarely small tree. Lamina texture thick, +/- leathery.
 ***Cercis occidentalis***
- 05 a Large (>1 cm) persisting stipules evident. ***Cercis chingii***
 b Stipules caducous. 06
- 06 a Lamina LS minutely to densely pubescent all over (10x LENS). 07
 b Leaflet LS (soon) glabrous, OR with minute pubescence restricted to veins/vein axils. ...10
- 07 a Lamina margin undulate. ***Cercis canadensis* var. *mexicana***
 b Lamina margin flat. 08
- 08 a Lamina LS basal veins 7-9. ***Cercis canadensis* subsp. *texensis***
 b Lamina LS basal veins 5-7. 09
- 09 a Lamina apex with acute acumen (10x LENS). ***Cercis racemosa***
 b Lamina apex with blunt acumen. ***Cercis yunnanensis***
- 10 a Lamina inequilateral, base obliquely cuneate; LS basal veins 3-5. ***Cercis chuniana***
 b Lamina equilateral, base (sub-)truncate to (deeply) cordate; LS basal veins 5-7(-9). 11
- 11 a Lamina LS basal veins 7-9. ***Cercis canadensis***
 b Lamina LS basal veins 5-7, OR 7. 12
- 12 a Lamina LS basal veins 5-7. ***Cercis glabra***
 b Lamina LS basal veins predominantly 7. 13
- 13 a Lamina LS basal vein axils predominantly minutely pubescent (10x LENS, examine
 several leaves). ***Cercis chinensis***
 b Lamina LS basal vein axils predominantly glabrous. **"*Cercis gigantea*"**

GROUP A2:

Foliage with phyllodes (sometimes juvenile bipinnate leaves may occur in part of the foliage).

01 a	Phyllode L/W ratio predominantly <4/1.	02
b	Phyllode L/W ratio >4/1.	08
02 a	Phyllode length <3 cm, <u>AND/OR</u> +/- triangular or obtriangular.	03
b	Phyllode length 4-10(-more) cm, elliptic to obovate.	04
03 a	Phyllode largest width clearly above the middle.	<i>Acacia pravissima</i>
b	Phyllode largest width in or below the middle.	<i>Acacia cultriformis</i>
04 a	Shoot, at least initially pubescent (10x LENS).	<i>Acacia podalyriifolia</i>
b	Shoot glabrous.	05
05 a	Phyllode with 2-4 major longitudinal veins.	<i>Acacia longifolia</i> subsp. <i>sophorae</i>
b	Phyllode with 1 longitudinal vein.	06
06 a	Phyllode length 2-6 cm.	<i>Acacia pataczekii</i>
b	Phyllode length 5-10(-more) cm.	07
07 a	Midvein eccentric.	<i>Acacia obliquinervia</i>
b	Midvein +/- central.	<i>Acacia pycnantha</i>
08 a	Phyllode with 1 longitudinal vein.	09
b	Phyllode with 2-4(-more) longitudinal veins.	13
09 a	Phyllode with 2-3 glands along margin (10x LENS).	<i>Acacia dodonaeifolia</i>
b	Phyllode with 1 gland, usually at/near base.	10
10 a	Bipinnate leaves often persist.	<i>Acacia rubida</i>
b	Bipinnate leaves usually absent.	11
11 a	Phyllode 10-25 cm long, basal gland prominent: flat and large (10x LENS).	<i>Acacia saligna</i>
b	Phyllode 6-16 cm long, basal gland not prominent/indiscernible.	12
12 a	Branchlet pruinose. Phyllode oblong elliptic, length <6 cm.	<i>Acacia covenyi</i>
b	Branchlet not pruinose. Phyllode oblanceolate, length 6-15 cm.	<i>Acacia retinodes</i>
13 a	Bipinnate leaves often persist.	<i>Acacia melanoxylon</i>
b	Bipinnate leaves usually absent.	14
14 a	Phyllode with 2-3 longitudinal veins.	15
b	Phyllode with 3-4(-more) longitudinal veins.	16

- 15 a Phyllode 5-20 cm, with apex soft. **Acacia longifolia**
 b Phyllode 1-5 cm, with apex pungent. **Acacia riceana**
- 16 a Full grown phyllode green. 17
 b Full grown phyllode grey-green or (sub-)glaucous. 18
- 17 a Phyllode length 5-8 cm, width 2-5 mm. **Acacia mucronata**
 b Phyllode length 10-20 cm, width 6-25(-35) mm. **Acacia maidenii**
- 18 a Phyllode length 4-12 cm, width 1-4 mm. **Acacia aneura**
 b Phyllode length 14-40 cm, width 2-7 mm. **Acacia stenophylla**

GROUP B1:
 Leaves trifoliolate.

- 01 a Terminal leaflet, in part of the foliage, small: midvein length ≤ 2 cm, AND longer than petiole (examine the whole plant). **+Laburnocytisus adamii**
 b Terminal leaflet midvein length clearly >2 cm, AND shorter than petiole. 02
- 02 a Long shoot completely pubescent with +/- appressed short hairs (10x LENS).
 **Laburnum anagyroides**
 b Long shoot (soon) glabrous, OR with scattered long hairs, OR variable (glabrous to pubescent in various parts of the foliage). 03
- 03 a Leaflet LS predominantly with scattered, long loose hairs along midvein (10 x LENS).
 **Laburnum alpinum**
 b Leaflet LS predominantly equally pubescent with +/- appressed short hairs, sometimes intermixed with long hairs. **Laburnum x watereri**

GROUP B2:
 Leaves pinnate AND/OR bipinnate.
 Leaflet margin (sometimes minutely!) crenulate (10x LENS).

- 01 a Leaves always pinnate (examine the whole foliage). 02
 b Leaves pinnate AND bipinnate. 04
- 02 a Petiolule ≤ 1 mm. Leaflet rhombic. **Gleditsia fera**
 b Petiolule 1-5 mm. Leaflets ovate or ovate-lanceolate to oblong. 03
- 03 a Leaflets 3-6 pairs. **Gleditsia macracantha**
 b Leaflets 4-8 pairs. **Gleditsia sinensis**

- 04 a Leaflet margin vaguely crenulate, appears entire at least in part of the foliage. Secondary veins rather anastomosing before reaching margin (10× LENS). *Gleditsia japonica*
- b Leaflet margin clearly crenulate with several secondary vein forks reaching margin. ... 05
- 05 a Leaflet LS midvein glabrous (10× LENS). *Gleditsia aquatica*
- b Leaflet LS midvein pubescent. 06
- 06 a Pinnae 4-7 pairs. Spines simple or trifid. *Gleditsia triacanthos*
- b Pinnae 3-4 pairs. Spines trifid to multi-forked. *Gleditsia caspica*

GROUP B3:
Leaves bipinnate (Leaflet margin entire).

- 01 a Upper pairs of pinnae with 4-7 pairs of ovate leaflets, AND the lower pairs are simple leaflets. *Gymnocladus dioica*
- b All pinnae with several pairs of leaflets, simple leaflets absent. 02
- 02 a Rachis armed with scattered decurved prickles. *Biancaea decapetala*
- b Rachis without decurved prickles. 03
- 03 a Pinnae in 1-6(-15) pairs + 1 terminal pinna (sometimes lacking). ... *Erythrostemon gilliesii*
- b Pinnae in pairs. 04
- 04 a Plant deciduous. 05
- b Plant evergreen (semi-evergreen in colder area). 06
- 05 a Pinnae in 4-12 pairs. Leaflet 8-12 mm. *Albizia julibrissin*
- b Pinnae in 3-6 pairs. Leaflet 15-40 mm. *Albizia kalkora*
- 06 a Branchlets clearly armed with stipular spines. Leaves often in fascicles. 07
- b Stipular spines absent. Leaves alternate. 08
- 07 a Stipular spines up to 2-3 cm. Branchlets minutely pubescent (10× LENS). *Vachellia caven*
- b Stipular spines up to 5-12(-more) cm. Branchlets glabrous. *Vachellia karroo*
- 08 a Phyllodes quite often present. *Acacia melanoxylon*
- b Phyllodes absent. 09
- 09 a Jugary gland and opening, at base of pinnae usually absent/indiscernible (10× LENS). .10
- b Jugary gland and opening, at base of several pinnae often prominent. 11
- 10 a Leaf small: 5 cm × 2,5 cm, petiole <1 cm. *Acacia cardiophylla*
- b Leaf large: 10-20(-more) cm, petiole 3-8 cm. *Paraserianthes lophantha*

- 11 a Pinnae in 2-4 pairs. *Acacia baileyana*
- b Pinnae in 6-20(-more) pairs. 12
- 12 a Emerging foliage glabrescent or with whitish-grey pubescence (10x LENS). 13
- b Emerging foliage with golden-yellow pubescence. 14
- 13 a Young shoots, rachis and leaves +/- minutely silvery-glaucous pubescent (10x LENS),
Pinnae closely spaced/touching. *Acacia dealbata*
- b Young shoots, rachis and leaves glabrous to puberulous. Pinnae widely spaced.
..... *Acacia decurrens*
- 14 a Jugary gland at base of each pair of pinnae (10x LENS). *Acacia parramattensis*
- b Jugary gland often absent in part of the pinnae. *Acacia irrorata*

GROUP B4:

Leaves pinnate (Leaflet margin entire).
Lateral buds concealed by leaf pulvini.

- 01 a Leaflet apex gradually acuminate, or acute to emarginate, BUT apiculus or bristle absent
(10x LENS). 02
- b Leaflet apex acute to emarginate AND apiculate to +/- bristly. 04
- 02 a Leaflet apex gradually acuminate. Terminal leaflet with 2 stipels, lateral leaflet with 1
stipel (10x LENS). *Platysprion platycarpum*
- b Leaflet apex emarginate or acute. Leaflets without stipels. 03
- 03 a Leaflets 9-13, closely set; +/- uniformly ovate oblong, AND apically +/- emarginate (10x
LENS). *Cladrastis delavayi*
- b Leaflets 5-11, distantly set; variable from elliptic to obovate or ovate (often on the same
leaf!), AND apically acute. *Cladrastis kentukea*
- 04 a Branchlets vivid green, AND green to olive-green for several successive years. 05
- b Branchlets often brownish to reddish, AND woody the second year. 06
- 05 a Leaflets 9-15, ovate to elliptic. *Styphnolobium japonicum*
- b Leaflets 13-19, elliptic. *Styphnolobium affine*
- 06 a Leaflets 7-13(-15), AND shoot +/- bristly pubescent to the naked eye. *Robinia hispida*
- b Leaflets 13-23. Shoot viscous and warty glandular, OR glabrous to slightly pubescent
(except inflorescence!). 07
- 07 a Shoot conspicuously viscous and warty glandular (10x LENS). *Robinia viscosa*
- b Shoot not warty glandular. 08
- 08 a Leaflet apically predominantly rounded and +/- emarginate. *Robinia pseudoacacia*
- b Leaflet apically predominantly (sub-)acute. *Robinia neomexicana*

GROUP B5:
Leaves pinnate (Leaflet margin entire).
Lateral buds visible.

01 a	Largest leaflet midvein length >3 cm.	02
b	Largest leaflet midvein length <3 cm.	08
02 a	Leaflet apex rounded to emarginate.	03
b	Leaflet apex acute.	04
03 a	Leaflets 4-10(-12).	Ceratonia siliqua
b	Leaflets 12(13)-24(25).	Tipuana tipu
04 a	Leaflets 19-25.	Sophora flavescens
b	Leaflets 7-15(-17).	05
05 a	Leaflets 7-11.	06
b	Leaflets 9-13, <u>OR</u> 9-15(-17).	07
06 a	Leaflet LS glabrous to glabrescent (10× LENS).	Maackia amurensis
b	Leaflet LS pubescent.	Maackia buergeri
07 a	Leaflets 9-13.	Maackia hupehensis
b	Leaflets 9-15(-17).	Maackia floribunda
08 a	Leaflets 7-25.	09
b	Leaflets 25-51(-more), at least so in part of the foliage.	13
09 a	Larger leaf length <10 cm.	10
b	Larger leaf length >10 cm.	11
10 a	Leaf length <3 cm.	Sophora prostrata
b	Leaf length 3-8 cm.	Sophora toromiro
11 a	Leaflet US glabrous or slightly pubescent in part (10× LENS).	Sophora howinsula
b	Leaflet US clearly appressed pubescent.	12
12 a	Leaflet margin clearly revolute (10× LENS).	Sophora macrocarpa
b	Leaflet margin rather flat.	Sophora tetraptera
13 a	Leaflet sessile, petiolule +/- absent/indiscernible (10× LENS).	Sophora fulvida
b	Leaflet with petiolule evident.	14
14 a	Leaflets all distant.	15
b	Leaflets closely set and (slightly) overlapping in part of the leaf.	16

- 15 a Leaflet largest width clearly above the middle. **Sophora longicarinata**
 b Leaflet largest width in the middle. **Sophora microphylla**
 16 a Leaflets rather overlapping in leaf apical part. **Sophora cassioides**
 b Leaflets rather overlapping in leaf basal part. **Sophora chathamica**

Taxa treated in this key.

Acacia aneura
Acacia baileyana
Acacia cardiophylla
Acacia covenyi
Acacia cultriformis
Acacia dealbata
Acacia decurrens
Acacia dodonaeifolia
Acacia irrorata
Acacia longifolia
 -subsp. *sophorae*
Acacia maidenii
Acacia melanoxylon
Acacia mucronata
Acacia obliquinervia
Acacia parramattensis
Acacia pataczekii
Acacia podalyriifolia
Acacia pravissima
Acacia pycnantha
Acacia retinodes
Acacia riceana
Acacia rubida
Acacia saligna
Acacia stenophylla
Albizia julibrissin
Albizia kalkora
Biancaea decapetala
Ceratonia siliqua
Cercis canadensis
 -var. *mexicana*
 -subsp. *texensis*
Cercis chinensis
Cercis chingii
Cercis chuniana
 "Cercis gigantea"
Cercis glabra
Cercis griffithii
Cercis occidentalis
Cercis racemosa
Cercis siliquastrum
Cercis yunnanensis
Cladrastis kentukea
Cladrastis delavayi

Erythrostemon gilliesii
Gleditsia aquatica
Gleditsia caspica
Gleditsia fera
Gleditsia japonica
Gleditsia macracantha
Gleditsia sinensis
Gleditsia triacanthos
Gymnocladus dioicus
 +*Laburnocytisus adamii*
 (Laburnum anagyroides + Cytisus purpureus)
Laburnum alpinum
Laburnum anagyroides
Laburnum × *watereri* (L.alpinum × L.anagyroides)
Maackia amurensis
Maackia buergeri
Maackia floribunda
Maackia hupehensis
Paraserianthes lophantha
Platyosprion platycarpum
Robinia hispida
Robinia neomexicana
Robinia pseudoacacia
Robinia viscosa
Sophora cassioides
Sophora chathamica
Sophora flavescens
Sophora fulvida
Sophora howinsula
Sophora longicarinata
Sophora macrocarpa
Sophora microphylla
Sophora prostrata
Sophora tetraptera
Sophora toromiro
Styphnolobium affine
Styphnolobium japonicum
Tipuana tipu
Vachellia caven
Vachellia karroo

Taxa referred to synonymy in this key (fide Plants of the World Online: [POWO](http://powo.science.kew.org/)).

Acacia caven → *Vachellia caven*
Acacia karroo → *Vachellia karroo*
Albizia lophantha → *Paraserianthes lophantha*
Caesalpinia decapetala → *Biancaea decapetala*
Caesalpinia gilliesii → *Erythrostemon gilliesii*
Caesalpinia japonica → *Biancaea decapetala*
Cercis reniformis → *Cercis canadensis* subsp. *texensis*
Cladrastis platycarpa → *Platyosprion platycarpum*

Cladrastis sinensis → *Cladrastis delavayi*
Gleditsia ferox → *Gleditsia triacanthos*
Gleditsia macracantha ~ *Gleditsia sinensis*
Gleditsia rolfei auct. → *Gleditsia fera*
Gymnocladus dioica → *Gymnocladus dioicus* (genus masculinum conservandum)
Sophora affine → *Styphnolobium affine*
Sophora japonica → *Styphnolobium japonicum*

